

Calciveyxol 38®

Solution for infusion

Good tissue tolerance with rapid effect

Composition:

Calcium gluconate	380 mg
(corresponding to Ca ²⁺ : 34mg or 0.85 mmol, respectively)	
Magnesium chloride hexahydrate	60mg
(corresponding to Mg ²⁺ : 7.2 mg or 0.30 mmol, respectively)	
Boric acid	50mg

Target species:

Cattle, sheep, pigs.

Indications:

Acute hypocalcaemic conditions.

As a supporting therapy for allergies, urticaria, haemorrhagic diathesis, uterine inertia

Recommended dose:

For slow intravenous application.

Cattle:

Acute hypocalcaemic conditions: 20 – 30 ml Calciveyxol® 38 per 50 kg b.w., intravenously (equivalent to 0.34 – 0.51 mmol Ca²⁺ and 0.12 – 0.18 mmol Mg²⁺ per kg b.w.)

Supportive therapy for allergies, urticaria, haemorrhagic diathesis, uterine atony: 15 – 20 ml Calciveyxol® 38 per 50 kg b.w., intravenously (equivalent to 0.26 – 0.34 mmol Ca²⁺ and 0.09 – 0.12 mmol Mg²⁺ per kg b.w.)

Sheep, calf, pig:

3 – 4 ml Calciveyxol® 38 per 10 kg b.w., intravenously (equivalent to 0.26 – 0.34 mmol Ca²⁺ and 0.09 – 0.12 mmol Mg²⁺ per kg b.w.)

Packages:

500 ml glass bottle

License holder:

Veyx Pharma GmbH, Germany

Importer and distributor:

Global Pharmaceutical Industries, Egypt